## BIDHAN CHANDRA KRISHI VISWAVIDYALAYA MEDIUM-RANGE WEATHER FORECAST AND AGRO-ADVISORY (Agromet Advisory Services, IMD, MoES)

Directorate of Research, B.C.K.V., Kalyani.



E-mail: <u>aaskalyani@gmail.com</u> bhowmickmahasweta2012@gmail.com

From Kalyani (Nadia, WB)

## Weather Forecast for Hooghly District

FORECAST PERIOD: : 19th May to 23th May, 2018

## Weather forecast for next five days

- ➤ Partly cloudy sky is expected during next few days .
- ➤ Light to medium rainfall is expected during next few days except 19th & 20th May & total will be 17 mm.
- ➤ Wind flow mainly from S E, S direction with 12-19 km/hr. speed during next 5 days is expected.
- Maximum relative humidity is expected to be around 75%-84% and minimum relative humidity is expected to be around 26% 38%.
- Maximum temperature is expected to be around  $33^{\circ}\text{C}$  - $36^{\circ}\text{C}$  and minimum temperature is expected to be around  $24^{\circ}\text{C}$  - $26^{\circ}\text{C}$ .

## Agrometeorological Advisory

Crop	Stage	Advisory
Kharif <b>M</b> aize	Sowing	Sowing of fodder crops may be done during next week. Adequate moisture should be maintained for maximum germination of seeds. Sowing should be done at a depth of 3-4 cm and at row to row spacing of 25-30 cm.
Legume vegetable	vegetative	For root rot apply (Carbendazim + Mancozeb) @2.0 g/ lit. of water as soil drenching. For aphid control of Lab Lab bean apply ash in case of lower infestation and incase of heavy infestation Thiomethoxam 0.3 SG @1g/lit. of water may be applied.
Boro Paddy	Panicle maturing stage	Do not need to irrigate the rice field as light rainfall has been forecasted . Don't harvest untill rice grains turn yellowish. Fully mature grains may be harvested in the morning hours.
Mango, Lichi	Fruit Setting	To prevent fruit drop apply NAA (Planofix) @ 3ml per 15 lt of water.
Jute	Seedling blight	In case of rotting of seedlings apply carbendazim 50 WP @1g/lit. of water or Cu-hydroxide 75 WP@ 2g/lit.of water