## BIDHAN CHANDRA KRISHI VISWAVIDYALAYA MEDIUM-RANGE WEATHER FORECAST AND AGRO-ADVISORY (Agromet Advisory Services, IMD, MoES)

Directorate of Research, B.C.K.V., Kalyani.



 $E\text{-mail: } \underline{aaskalyani@gmail.com} \ , \\ bhowmickmahasweta 2012@gmail.com$ 

From Kalyani (Nadia, WB)

## Weather Forecast for Nadia District

FORECAST PERIOD: : 12th May to 16th May, 2018

## Weather forecast for next five day

- $\blacktriangleright$  Partly cloudy sky is expected during next few days expect 12th & 13th May(clear sky).
- ➤ Light rainfall is expected during next few days & total will be 53 mm.
- ➤ Wind flow mainly from E, S and SE direction with 6-14 km/hr. speed during next 5 days is expected.
- Maximum relative humidity is expected to be around 77%-86% and minimum relative humidity is expected to be around 31% 45%.
- ➤ Maximum temperature is expected to be around 32°C 35°C and minimum temperature is expected to be around 24 °C -25°C.

## **Agrometeorological Advisory**

| Crop                      | Stage                        | Advisory  |
|---------------------------|------------------------------|---|
| Kharif<br><b>Maize</b>    | Sowing                       | Sowing of fodder crops may be done during next week. Adequate moisture should be maintained for maximum germination of seeds. Sowing should be done at a depth of 3-4 cm and at row to row spacing of 25-30 cm. |
| Green<br>manuring<br>crop | Sowing                       | Sowing of green manuring crop like cluster bean, dhaincha etc may be advised where harvesting of rabi crop is over.  Ensure proper moisture for proper germination and growth of crop.                          |
| Boro<br>Paddy             | Panicle<br>maturing<br>stage | Do not need to irrigate the rice field as light rainfall has been forecasted . Don't harvest untill rice grains turn yellowish. Fully mature grains may be harvested in the morning hours.                      |
| Mango,<br>Lichi           | Fruit Setting                | To prevent fruit drop apply NAA (Planofix) @ 3ml per 15 lt of water.  |
| Jute                      | Seedling                     | Thinning maybe done and maintain the plant population in the main field.  |