



Gramin Krishi Mausam Sewa

District Level Agromet Advisory Bulletin
Bidhan Chandra Krishi Viswavidyalaya
 Mohanpur, Dist- Nadia, West Bengal



Agromet Advisory Bulletin

Date : 23-11-2021

Weather Forecast of PASCHIM-BARDHAMAN(West Bengal) Issued On : 2021-11-23(Valid Till 08:30 IST of the next 5 days)

Parameter	2021-11-24	2021-11-25	2021-11-26	2021-11-27	2021-11-28
Rainfall(mm)	0.0	0.0	0.0	0.0	0.0
Tmax(°C)	30.0	29.0	28.0	28.0	27.0
Tmin(°C)	17.0	16.0	16.0	15.0	15.0
RH-I(%)	36	48	54	67	60
RH-II(%)	26	26	33	28	30
Wind Speed(kmph)	8.9	7.0	6.4	6.6	7.9
Wind Direction(Degree)	349	22	25	26	345
Cloud cover(octa)	0	0	0	0	1

Weather Summary/Alert:

No rainfall is likely to occur during next few days. Clear sky is expected on next few days . Wind speed is expected to be 6-9 km / hour. Max. RH is expected to be 36% to 67%. Maximum temperature is expected to be around 27.0-30.0 deg C and minimum temperature is expected to be around 15.0-17.0 deg C.

General Advisory:

No rain is expected , spraying , fertilizer application , harvesting may be done .

SMS Advisory:

Please download Mausam , Meghdoot & Damini App at your smart phone to get weather forecast and agromet advisory .

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	Continue rice harvesting . After drying & threshing ,grains should be properly dried to right moisture level. Don't burn the residue of the crop.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
-------------------------	--------------------------------

Horticulture(Varieties)	Horticulture Specific Advisory
POTATO	<p>Complete planting of potato in between 15th to 24th Nov. N:P:K dose 150 : 150 : 150 kg / ha may give the good result . The crop matures in 80-90 days. Medium high land , sandy loam to loamy soil , slightly acidic to neutral ph is ideal for potato cultivation. One deep ploughing , 2-3 cross & shallow ploughing make the soil well pulverized , loose , & friable with good tilth . 10-15 ton well rotten organic manure must be mixed during first ploughing . Apply phorate 10 G @ 15 Kg/ha to control soil borne insect. Collect certified seed tuber from Govt. approved organization . 25-45 g seed tuber must be treated with trichoderma viridi or pseudomonas fluorescence solution. Recommended dose of N :P:K is 180-200 : 150 : 150 Kg/ha. Nitrogen should be applied in split doses . Zn & B may be applied in case of micronutrient deficiency in the form of ZnSo₄ (25 Kg/ha) and Borax (10 Kg/ha) at the time of final land preparation .</p>