4/8/22, 4:28 PM District Advisory



Gramin Krishi Mausam Sewa District Level Agromet Advisory Bulletin Bidhan Chandra Krishi Viswavidyalaya



Mohanpur, Dist- Nadia, West Bengal

Agromet Advisory Bulletin

Date: 08-04-2022

Weather Forecast of District HOOGHLY(West Bengal) Issued On: 2022-04-08(Valid Till 08:30 IST of the next 5 days)

| Parameter | 2022-04-09 | 2022-04-10 | 2022-04-11 | 2022-04-12 | 2022-04-13 |
|------------------------|------------|------------|------------|------------|------------|
| Rainfall(mm) | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Tmax(°C) | 34.0 | 34.0 | 33.0 | 33.0 | 34.0 |
| Tmin(°C) | 25.0 | 26.0 | 26.0 | 26.0 | 27.0 |
| RH-I(%) | 74 | 82 | 78 | 83 | 83 |
| RH-II(%) | 14 | 20 | 39 | 33 | 38 |
| Wind Speed(kmph) | 23.3 | 24.6 | 25.8 | 22.1 | 21.4 |
| Wind Direction(Degree) | 201 | 201 | 201 | 167 | 167 |
| Cloud Cover(Octa) | 0 | 0 | 0 | 1 | 2 |

Weather Summary/Alert:

No rainfall is likely to occur during next few days. Clear sky is expected on next few days. Wind speed is expected to be 21-26 km / hour. Max. RH is expected to be 74% to 83%. Maximum temperature is expected to be around 33.0-34.0 deg C and minimum temperature is expected to be around 25.0-27.0 deg C.

General Advisory:

No rainfall is expected in the next few days. Spraying and harvesting may be done.

SMS Advisory:

No rainfall is expected in the next few days. Spraying and harvesting may be done.

Horticulture Specific Advisory:

| Horticulture(Varieties) | Horticulture Specific Advisory |
|-------------------------|---|
| BRINJAL | Fruit Setting Stage: Brinjal fruit & shoot borer may be seen at this stage. As soon as the insect is detected, the affected parts should be clipped along with the insect and destroyed. Fruits showing any boring should be picked and destroyed. Weed control is must. Spray with Profenophos + Cypermethrin 1.5ml + Acitamiprid 1gm at 15 days interval. After spraying apply prime pack 1.0-2.6-2.6 liquid fertilizer (5 ml) + Humax 1 ml/lt + 10-26-26 (10 kg/ bigha) + I kg Sagarika dana +5 kg Neemcake as topdressing. One solar light trap per bigha will be useful. |
| | Chilli • Fruiting stage • Mite and thrips attack. For mite attack spray Dicofol or Propergite . For thrips attack spray Thiamithoxam . |