



Gramin Krishi Mausam Sewa
District Level Agromet Advisory Bulletin
Bidhan Chandra Krishi Viswavidyalaya
 Mohanpur, Dist- Nadia, West Bengal



Agromet Advisory Bulletin

Date : 06-05-2022

Weather Forecast of District HOWRAH(West Bengal) Issued On : 2022-05-06(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-05-07	2022-05-08	2022-05-09	2022-05-10	2022-05-11
Rainfall(mm)	4.0	1.0	1.0	5.0	10.0
Tmax(°C)	36.0	36.0	36.0	34.0	33.0
Tmin(°C)	26.0	26.0	27.0	26.0	25.0
RH-I(%)	72	83	77	80	80
RH-II(%)	31	31	36	36	38
Wind Speed(kmph)	12.5	15.1	17.2	20.0	18.7
Wind Direction(Degree)	193	180	162	159	159
Cloud Cover(Octa)	1	1	2	0	0

Weather Summary/Alert:

Light to no rainfall is likely to occur during next few days. Clear to partly cloudy sky is expected on next few days . Wind speed is expected to be 12-20 km / hour. Max. RH is expected to be 72% to 80 % . Maximum temperature is expected to be around 33.0-36.0 deg C and minimum temperature is expected to be around 25.0-27.0 deg C.

General Advisory:

Spraying ,fertilizer application and harvesting maybe avoided because rainfall is expected on next fewdays.

SMS Advisory:

Spraying ,fertilizer application and harvesting maybe avoided because rainfall is expected on next fewdays.

Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
DAINCHA	Sowing of green manuring crop like cluster bean, dhaincha etc may be advised where harvesting of rabi crop is over .Ensure proper moisture for proper germination and growth of crop.

Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
-------------------------	--------------------------------

Horticulture(Varieties)	Horticulture Specific Advisory
BRINJAL	Fruit Setting Stage : Brinjal fruit & shoot borer may be seen at this stage . As soon as the insect is detected , the affected parts should be clipped along with the insect and destroyed .Fruits showing any boring should be picked and destroyed . Weed control is must . Spray with Profenophos + Cypermethrin 1.5ml + Acitamiiprid 1gm at 15 days interval. After spraying apply prime pack 1.0-2.6-2.6 liquid fertilizer (5 ml) + Humax 1 ml/lit + 10-26-26 (10 kg/ bigha) + 1 kg Sagarika dana +5 kg Neemcake as topdressing . One solar light trap per bigha will be useful .
CHILLI	Chilli • Fruiting stage • Mite and thrips attack. For mite attack spray Dicofol or Propergite . For thrips attack spray Thiamithoxam .
BANANA	In the month of May, to counteract storm , tie each banana plant against a bamboo with a strong rope and save your plant from lodging.