



**Gramin Krishi Mausam Sewa**  
**District Level Agromet Advisory Bulletin**  
**Bidhan Chandra Krishi Viswavidyalaya**  
Mohanpur, Dist- Nadia, West Bengal



## Agromet Advisory Bulletin

Date : 15-12-2023

Weather Forecast of District MEDINIPUR-EAST(West Bengal) Issued On : 2023-12-15(Valid Till 08:30 IST of the next 5 days)

Parameter	2023-12-16	2023-12-17	2023-12-18	2023-12-19	2023-12-20
Rainfall(mm)	0.0	0.0	0.0	0.0	0.0
Tmax(°C)	27.0	27.0	27.0	27.0	26.0
Tmin(°C)	12.0	12.0	13.0	13.0	12.0
RH-I(%)	47	43	42	44	48
RH-II(%)	27	26	27	28	24
Wind Speed(kmph)	11	11	10	11	14
Wind Direction(Degree)	300	15	15	285	338
Cloud Cover(Octa)	0	0	1	2	3

### Weather Summary/Alert:

No rainfall is predicted on next 5 days. • Sky will be mainly clear in next 5 days . • Wind speed will be 11-15 km/hr and the predominant wind direction will be Northerly to Westerly . • Maximum temperature is expected to be around 26.0-28.0 degree and minimum temperature is likely to be 12.0-13.0degree. • Maximum and minimum relative humidity will be in the range of 42-48% and 24-28%.

### General Advisory:

As per extended range forecast no rainfall is recorded from 23.11.23 to 29.11.23 and normal rainfall is recorded from 30.11.23 to 6.12.23. Based on MME forecast above normal rainfall is predicted from 8.12.23 to 21.12.23. Based on MME forecast Tmax will be below normal in Week 1 and 2. and for Tmin, it will show below normal in Week 1 and week 2.

### SMS Advisory:

As minimum temperature is forecasted below normal, apply dry food for poultry and fishery only in day time for avoiding low temperature.

### Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	Boro paddy Sowing • Prepare low land field for nursery bed preparation in an area of 1/10th of total main field. • The bed should be 1mt in width and separate the seed bed by 20 cm depth irrigation channel. • Apply basal fertilizer viz. urea@750g/katha, SSP@2kg/katha and MOP@500g/katha. • Seeds should be treated with Bavistin@2g/kg of seed or Thirum 70WP@2.5g/kg of seed for • dry seed treatment. For wet seed treatment seeds should be dipped for 8-10 hours by • MEMC@15g/liter of water and then keep in gunny bag for 40-

Crop(Varieties)	Crop Specific Advisory
	50hours for better germination. Boro paddy For early sown paddy Seedling stage chance of Root rot and yellowing of seedlings due to low temperature • Make shade by transparent polythene over seed bed • Follow water treatment (fill the bed with water during evening and remove the water at morning hours ) for temperature conservation when seedlings are already grown up • Apply boron and zinc EDTA 0.5g/liter of water for getting healthy seedlings • Apply Diethane M-45@2.5g/liter of water after complete drain out of water from seed bed with growing seed lings • Avoid urea application on growing seed bed for avoiding its cooling effect.

**Horticulture Specific Advisory:**

Horticulture(Varieties)	Horticulture Specific Advisory
CAULIFLOWER	Winter vegetables (Cabbage, Cauliflower) now in head formation stage • Infestation of spodoptera may transform from harvested aman paddy field • Apply Spinosad @1.5ml/liter of water

**Live Stock Specific Advisory:**

Live Stock(Varieties)	Live Stock Specific Advisory
COW	Cattle All stages • Take special care for newly delivered mother cow • Keep clean and dry the new born calf • Cover the openings of shed by jute bag • Feed the cattle dry food • Cover the young one by jute bag