



Gramin Krishi Mausam Sewa
District Level Agromet Advisory Bulletin
Bidhan Chandra Krishi Viswavidyalaya
 Mohanpur, Dist- Nadia, West Bengal



Agromet Advisory Bulletin

Date : 09-02-2024

Weather Forecast of District MEDINIPUR-EAST(West Bengal) Issued On : 2024-02-09(Valid Till 08:30 IST of the next 5 days)

Parameter	2024-02-10	2024-02-11	2024-02-12	2024-02-13	2024-02-14
Rainfall(mm)	0.0	0.0	0.0	0.0	0.0
Tmax(°C)	26.0	26.0	27.0	28.0	29.0
Tmin(°C)	13.0	14.0	15.0	16.0	18.0
RH-I(%)	39	41	42	46	57
RH-II(%)	22	24	27	25	27
Wind Speed(kmph)	11	9	6	8	10
Wind Direction(Degree)	30	342	288	207	180
Cloud Cover(Octa)	1	0	3	7	5

Weather Summary/Alert:

No rainfall is predicted in next 5 days. • Sky will be mainly clear to partly cloudy in next 5 days . • Wind speed will be 6-11 km/hr and the predominant wind direction will be Northerly. • Maximum temperature is expected to be around 26.0-29.0 degree and minimum temperature is likely to be 13.0-18.0 degree. • Maximum and minimum relative humidity will be in the range of 39-57% and 22-27%.

General Advisory:

As per extended range forecast large excess rainfall is recorded from 25.1.24 to 7.2.24. Based on MME forecast below normal rainfall is predicted from 9.2.24 to 22.2.24 . Based on MME forecast Tmax will be below normal in Week 1 and week 2. For Tmin, it will show below normal in Week 1 and week 2

SMS Advisory:

No rainfall is predicted. Lathyrus now in flowering stage, apply 2% DAP for better flowering.

Crop Specific Advisory:

Crop (Stage)	Crop Specific Advisory
RICE (Vegetative)	Boro paddy --Transplanting almost complete /vegetative stage • Where seedlings are at 30-35 days old, there only transplanting can be continued. • Apply pre emergence weedicide just after 3-4 days of transplanting. Early transplanted boro paddy --Blast attack --Vegetative stage • Where transplanting completed 10-15 days ago apply chlorothalonipol (fatera) for preventing stem borer attack. It prevent the stem borer attack. • Complete first

Crop (Stage)	Crop Specific Advisory
	<p>top dressing with urea @14kg per bigha • Before top dressing drain out water from field then apply fertilizer and after proper puddling and wait for 24 hours again enter the water in the field. • Those who did not apply Tricyclozole in seed bed before transplanting apply Tricyclozole @ 5-6g per 10 liter of water as a preventive method. • For fungicide application first drain out water from field then keep the field dry and spray any fungicide and then after 48 hours again enter the water in the field. • For better result follow community spraying.</p>

Horticulture Specific Advisory:

Horticulture (Stage)	Horticulture Specific Advisory
CHILLI (Flowering)	<p>Chilli --Vegetative to flowering stage --Leaf curl due to strong Northern wind. • Spray Imidachlorpit @3ml per 10 liter of wate or Thiamethoxon @ 1g per liter of water.</p>
BRINJAL (Vegetative)	<p>Brinjal --Vegetative stage --Shoot borer attack • Apply Spinosad@1.5ml per liter of water during afternoon time. • No spraying should be done during morning hours in huge foggy weather.</p>