2/16/24, 2:42 PM District Advisory



Gramin Krishi Mausam Sewa District Level Agromet Advisory Bulletin Bidhan Chandra Krishi Viswavidyalaya



Mohanpur, Dist- Nadia, West Bengal

Agromet Advisory Bulletin

Date: 16-02-2024

Weather Forecast of District NADIA(West Bengal) Issued On: 2024-02-16(Valid Till 08:30 IST of the next 5 days)

Parameter	2024-02-17	2024-02-18	2024-02-19	2024-02-20	2024-02-21
Rainfall(mm)	0.0	0.0	0.0	0.0	0.0
Tmax(°C)	28.0	28.0	29.0	30.0	30.0
Tmin(°C)	14.0	14.0	15.0	17.0	19.0
RH-I(%)	51	46	41	68	64
RH-II(%)	23	21	18	22	25
Wind Speed(kmph)	10	10	6	8	13
Wind Direction(Degree)	326	297	292	162	206
Cloud Cover(Octa)	0	0	0	1	1

Weather Summary/Alert:

No rainfall is expected on next few days. Clear sky is expected in the next few days. Wind speed is expected 6-9 km / hour. Max. RH is expected to be 41% to 68%. Maximum temperature is expected to be around 28.0-30.0 deg C and minimum temperature is expected to be around 14.0-19.0 deg C.

General Advisory:

Please download Meghdoot app at your smart phone to get medium range weather forecast and agromet advisory.

SMS Advisory:

Spraying, fertilizer application and harvesting may be done because no rainfall is expected.

Crop Specific Advisory:

Crop (Stage)	Crop Specific Advisory
RICE (Vegetative)	Boro Paddy: Transplanting: Complete transplanting where seedlings are 35-40 days old. During main field preparation apply compost and basal dose of fertilizer as required. Light rainfall of last few days may cause fungal infection. Apply Trycyclozole @ 0.5 g per lt of water and any pesticide as preventive measure.l

Horticulture Specific Advisory:

Horticulture (Stage)	Horticulture Specific Advisory
	Late vegetative to tuberization stage: Micronutients may be applied at this stage to enhance the growth and quality of tuber.
CABBAGE (Vegetative)	Present weather is suitable for cultivation of winter vegetables .Leaf cutting caterpillar can be collected and killed by hand. Spraying may be done as

2/16/24, 2:42 PM District Advisory

Horticulture (Stage)	Horticulture Specific Advisory			
	required. To control White fly alternate spraying of Difenthiuron and			
	Azoxystrobin + Tebuconazole is recommended.			