2/20/24, 9:00 PM District Advisory



# Gramin Krishi Mausam Sewa District Level Agromet Advisory Bulletin Bidhan Chandra Krishi Viswavidyalaya



Mohanpur, Dist- Nadia, West Bengal

## **Agromet Advisory Bulletin**

Date: 20-02-2024

Weather Forecast of District HOOGHLY(West Bengal) Issued On: 2024-02-20(Valid Till 08:30 IST of the next 5 days)

Parameter	2024-02-21	2024-02-22	2024-02-23	2024-02-24	2024-02-25
Rainfall(mm)	0.0	0.0	5.0	2.0	2.0
Tmax(°C)	31.0	32.0	32.0	31.0	30.0
Tmin(°C)	22.0	22.0	22.0	21.0	21.0
RH-I(%)	73	64	76	63	48
RH-II(%)	25	24	23	35	30
Wind Speed(kmph)	10	12	15	15	13
Wind Direction(Degree)	202	189	225	0	25
Cloud Cover(Octa)	2	4	3	1	1

### Weather Summary/Alert:

No rainfall to light rainfall (22.2.24) is expected on next few days. Partly cloudy to clear sky is expected in the next few days. Max. RH is expected to be 48% to 76%. Maximum temperature is expected to be around 30.0-32.0 deg C and minimum temperature is expected to be around 21.0-22.0 deg C.

#### **General Advisory:**

Please download Meghdoot app at your smart phone to get medium range weather forecast and agromet advisory.

### **SMS Advisory:**

Spraying, fertilizer application and harvesting may be restricted on 22.2.24 because light rainfall is expected.

#### **Crop Specific Advisory:**

Crop (Stage)	Crop Specific Advisory		
RICE (Vegetative )	Boro Paddy: Transplanting: Complete transplanting where seedlings are 35-40 days old. During main field preparation apply compost and basal dose of fertilizer as required. Light rainfall of last few days may cause fungal infection. Apply Trycyclozole @ 0.5 g per lt of water and any pesticide as preventive measure. l		

### **Horticulture Specific Advisory:**

2/20/24, 9:00 PM District Advisory

Horticulture (Stage)	Horticulture Specific Advisory		
	Late vegetative to tuberization stage: Micronutients may be applied at this stage to enhance the growth and quality of tuber.		
CABBAGE (Vegetative)	Present weather is suitable for cultivation of winter vegetables .Leaf cutting caterpillar can be collected and killed by hand. Spraying may be done as required . To control White fly alternate spraying of Difenthiuron and Azoxystrobin + Tebuconazole is recommended.		