



Gramin Krishi Mausam Sewa
District Level Agromet Advisory Bulletin
Bidhan Chandra Krishi Viswavidyalaya
 Mohanpur, Dist- Nadia, West Bengal



Agromet Advisory Bulletin

Date : 20-02-2024

Weather Forecast of District HOWRAH(West Bengal) Issued On : 2024-02-20(Valid Till 08:30 IST of the next 5 days)

Parameter	2024-02-21	2024-02-22	2024-02-23	2024-02-24	2024-02-25
Rainfall(mm)	0.0	0.0	4.0	1.0	4.0
Tmax(°C)	31.0	32.0	32.0	31.0	30.0
Tmin(°C)	22.0	22.0	22.0	21.0	21.0
RH-I(%)	71	70	70	67	56
RH-II(%)	26	25	25	34	32
Wind Speed(kmph)	13	18	19	14	12
Wind Direction(Degree)	214	202	216	11	30
Cloud Cover(Octa)	1	3	2	1	1

Weather Summary/Alert:

No rainfall to light rainfall (22.2.24, 24.2.24) is expected on next few days. Partly cloudy to clear sky is expected in the next few days. Max. RH is expected to be 48% to 76%.Maximum temperature is expected to be around 30.0-32.0 deg C and minimum temperature is expected to be around 21.0-22.0 deg C.

General Advisory:

Please download Meghdoot app at your smart phone to get medium range weather forecast and agromet advisory.

SMS Advisory:

Spraying , fertilizer application and harvesting may be restricted on 22.2.24 because light rainfall is expected .

Crop Specific Advisory:

Crop (Stage)	Crop Specific Advisory
RICE (Vegetative)	Boro Paddy : Transplanting : Complete transplanting where seedlings are 35-40 days old. During main field preparation apply compost and basal dose of fertilizer as required . Light rainfall of last few days may cause fungal infection . Apply Tricyclozole @ 0.5 g per lt of water and any pesticide as preventive measure .l

Horticulture Specific Advisory:

Horticulture (Stage)	Horticulture Specific Advisory
POTATO (Vegetative)	Late vegetative to tuberization stage : Micronutrients may be applied at this stage to enhance the growth and quality of tuber .
CABBAGE (Vegetative)	Present weather is suitable for cultivation of winter vegetables .Leaf cutting caterpillar can be collected and killed by hand. Spraying may be done as required . To control White fly alternate spraying of Difenthiuron and Azoxystrobin + Tebuconazole is recommended.